
CERAMENT[®] G




Article number EU: 5ml A0535-02 EU: 10ml A0535-01

US: 5ml A0535-06 US: 10ml A0535-05





- Remove the cap from the mixing liquid.
- Keep the vial stable.
- Push one of the spikes through the membrane of the vial.



- Attach the purple syringe to the spike.
- Withdraw saline only up to the red mark on the syringe, there will be saline left in the glass vial.
- Disconnect the saline-filled syringe from the spike.




- Push the second spike through the membrane of the gentamicin vial.
- Attach the saline-filled syringe to the spike.




- Transfer all the liquid to the gentamicin vial.
- Dissolve all the gentamicin powder.




Withdraw all the gentamicin solution back into the syringe. This may require the spike to be slightly backed out.



Retract the blue handle and remove the red plunger stop ring.







- b) Attach valve, clear end to the powder-filled syringe, and turn clockwise.
- c) Attach syringe with gentamicin solution to blue end of valve, press and turn clockwise.




- Inject all the gentamicin solution into the syringe with powder.
- Keep the plunger pushed to the bottom to avoid back flush.
- Detach the syringe from the valve.



- After the liquid has been injected, start the stopwatch immediately.
- Mix with rotation for 30 seconds, at 1 complete stroke per second.
- Push the handle all the way down and pull it all the way up, and rotate at its end positions.



- Fully retract blue handle.
- Lock plunger by turning blue collar clockwise to click.



CERAN

- Attach dispensing syringe to valve.
- Immediately transfer paste with numbers facing user. When full, paste will begin to ooze from under sleeve. Stop filling when this occurs.



Remove the red stopper and the paste is now ready to use.







PR 01326-02 en AU EU ROW US 01-2024